

## VU Research Portal

### Improving Walking Capacity by Surgical Correction of Equinovarus Foot Deformity in Adult Patients with Stroke Or Traumatic Brain Injury: A Systematic Review

Renzenbrink, G.J.; Buurke, J.H.; Nene, A.V.; Geurts, A.C.H.; Kwakkel, G.; Rietman, J.S.

***published in***

Journal of Rehabilitation Medicine  
2012

***DOI (link to publisher)***

[10.2340/16501977-1012](https://doi.org/10.2340/16501977-1012)

***document version***

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

***citation for published version (APA)***

Renzenbrink, G. J., Buurke, J. H., Nene, A. V., Geurts, A. C. H., Kwakkel, G., & Rietman, J. S. (2012). Improving Walking Capacity by Surgical Correction of Equinovarus Foot Deformity in Adult Patients with Stroke Or Traumatic Brain Injury: A Systematic Review. *Journal of Rehabilitation Medicine*, 44(8), 614-623.  
<https://doi.org/10.2340/16501977-1012>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

## REVIEW ARTICLE

# IMPROVING WALKING CAPACITY BY SURGICAL CORRECTION OF EQUINOVARUS FOOT DEFORMITY IN ADULT PATIENTS WITH STROKE OR TRAUMATIC BRAIN INJURY: A SYSTEMATIC REVIEW

Gerbert J. Renzenbrink, MD<sup>1,2</sup>, Jaap H. Buurke, PT, PhD<sup>1,2</sup>, Anand V. Nene, MD, PhD<sup>2</sup>, Alexander C. H. Geurts, MD, PhD<sup>3,4</sup>, Gert Kwakkel, PhD<sup>5,6</sup> and Johan S. Rietman, MD, PhD<sup>2,7</sup>

From the <sup>1</sup>Roessingh Rehabilitation Center, Stroke Unit, <sup>2</sup>Roessingh Research and Development, Enschede, <sup>3</sup>Radboud University Nijmegen, Medical Center, Department of Rehabilitation, <sup>4</sup>St Maartenskliniek Research, Development and Education, Nijmegen, The Netherlands, <sup>5</sup>Vrije Universiteit Medical Center Amsterdam, Department of Rehabilitation, Research institute MOVE, <sup>6</sup>University Medical Center Utrecht, Department of Rehabilitation Medicine and Sports, Rudolf Magnus Institute and <sup>7</sup>Laboratory of Biomechanical Engineering, University of Twente, Enschede, The Netherlands

**Objective:** Equinovarus foot deformity following stroke or traumatic brain injury compromises walking capacity, interfering with activities of daily living. In soft-tissue surgery the imbalanced muscles responsible for the deviant position of the ankle and foot are lengthened, released and/or transferred. However, knowledge about the effectiveness of surgical correction is limited. The aim of the present study was to carry out a systematic review of the literature to assess the effects of surgical correction of equinovarus foot deformity in patients with stroke or traumatic brain injury.

**Methods:** A systematic search of full-length articles in the English, German or Dutch languages published from 1965 to March 2011 was performed in PubMed, EMBASE, CINAHL, Cochrane and CIRRIE. The identified studies were analysed following the International Classification of Functioning, Disability and Health criteria.

**Results:** A total of 15 case series, case control and historically controlled studies (CEBM level 4) were identified, suggesting that surgical correction of equinovarus foot deformity is a safe procedure that is effective in terms of re-obtaining a balanced foot position, improving walking capacity and diminishing the need for orthotic use.

**Discussion:** Further validation of surgical correction of equinovarus foot deformity following stroke or traumatic brain injury is required, using higher level study designs with validated assessment tools. Comparing surgical techniques with other interventions is necessary to generate evidence upon which treatment algorithms could be based.

**Key words:** equinovarus; foot deformity; stroke; traumatic brain injury; orthopaedic surgery; adult.

J Rehabil Med 2012; 44: 614–623

Correspondence address: Gerbert J. Renzenbrink, Roessingh Rehabilitation Center, Roessinghbleekweg 33, NL-7522 AH Enschede, The Netherlands. E-mail: b.renzenbrink@roessingh.nl

Submitted December 21, 2011; accepted May 4, 2012

## INTRODUCTION

Patients with an upper motor neuron lesion, such as in stroke and traumatic brain injury (TBI), often experience disruption

of the functional balance between agonistic and antagonistic muscle activity. In the affected lower extremity, such muscular imbalance often causes deformities of the ankle and foot. Although different types of acquired ankle and foot deformity following stroke and TBI have been described, equinovarus deformity is most characteristic and most frequently seen (1, 2). In the lower limb, involuntary activity of the plantar flexors and invertors of the ankle and foot, combined with paresis of the dorsal flexors and evertors, may explain the dynamics in the development of equinovarus foot deformity (3, 4).

Impaired walking capacity is a well-known consequence of stroke and TBI, and regaining independent gait is considered to be a primary goal in the rehabilitation of these patients (5, 6). Previous prospective cohort studies have shown that 60–80% of stroke survivors are able to walk independently at 6 months post-stroke (7–9). Available data, albeit limited, regarding walking capacity have shown that up to 70% of TBI survivors will also re-gain independent gait within the first 6 months (6, 10). Besides the fact that independent gait is highly related to independence in activities of daily living (ADL) (5), a number of studies have shown that the extent of gait recovery also differentiates patients who are housebound from those who are unlimited community walkers (11–13).

Equinovarus foot deformity compromises several prerequisites of walking (14). It interferes with foot clearance in the swing phase, with appropriate prepositioning of the foot at the end of the swing phase, with loading of the stance leg and with ankle stability (and postural balance) during the stance phase. As a result, these patients experience an increased risk of falling and are frequently unable to walk either unassisted or without orthotic devices (15).

A number of non-invasive and invasive options to treat equinovarus foot deformity following stroke or TBI have been proposed in the literature. These treatment options include orthotics (16), chemical denervation (17–19), neurosurgical denervation (20), functional electrical stimulation (21, 22) and soft-tissue surgery. All these interventions aim to correct the deviant position of the ankle and foot.

In soft-tissue surgery the imbalanced muscles responsible for plantar flexion and inversion of the ankle and foot are length-

ened, released and/or transferred. As a result, the muscle forces that act on the ankle–foot complex are balanced and equinovarus deformity is corrected (23, 24). However, in evidence-based rehabilitation guidelines the surgical correction of equinovarus deformity is not commonly addressed (25, 26). Indeed, in the average rehabilitation setting, ankle–foot orthoses and orthopaedic footwear often constitute the first choice of treatment. It is an important question whether this preference is based on convenience or on clinical evidence (27).

To our knowledge the effect of surgical correction of equinovarus foot deformity following stroke or TBI is indistinct. In addition, the criteria for selecting patients for soft-tissue surgery are heterogeneous, hindering optimal clinical decision-making (28).

Hence, the aim of the present study was to carry out a systematic review of the literature to assess the effects of surgical correction of equinovarus foot deformity in patients with stroke or TBI. Subsequently, recommendations with respect to selecting patients with equinovarus foot deformity are given to support the clinical decision-making process in patients with equinovarus foot deformity who are eligible for surgical correction.

## METHODS

### Definitions

- *Stroke* is defined as “an acute neurological dysfunction of vascular origin with sudden (within seconds) or at least rapid (within hours) occurrence of symptoms and signs corresponding to the involvement of focal areas of the brain” (29).
- *Traumatic brain injury* is defined as “damage to brain tissue caused by an external mechanical force as evidenced by medically documented loss of consciousness or post-traumatic amnesia due to brain trauma or by objective neurological findings that can be reasonably attributed to TBI on physical examination or mental status examination” (30).
- *Equinovarus foot deformity* is defined as “a combination of a plantar-flexed, inverted and adducted foot, (either dynamic or structural), acquired after stroke or TBI with possibly various degrees of severity of the different components”.
- *Walking capacity* is defined as “the degree of autonomy in walking, with or without the aid of appropriate assistive devices (such as canes or walkers), safely and sufficiently to carry out mobility-related activities of daily living”.
- *Capacity*, according to the World Health Organization (WHO) International Classification of Functioning Disability and Health (ICF), is a “qualifier” that describes a patient’s ability to execute a task or action, but *does not* qualify what a patient does in his or her current environment (performance) (31).

### Study identification

A systematic search was conducted of articles published in PubMed, EMBASE, CINAHL, Cochrane Controlled Trials (Rehabilitation and Related Therapies), Center for International Rehabilitation Research Information and Exchange (CIRRIE, <http://cirrie.buffalo.edu/>), and National Rehabilitation Information Center for Independence REHABDATA (<http://www.naric.com>) databases between 1965 and March 2011. REHABDATA was consulted for rehabilitation research conducted within the USA. CIRRIE includes research from all areas of rehabilitation conducted outside the USA, starting in 1990. The following MeSH headings and key words were used: “equinus, equinovarus, foot deformity, foot deformities, hemiplegia, hemiparesis, stroke, cerebrovascular disorders, head injury, traumatic brain injury, upper

motor neuron, orthopedics, neurorehabilitation, surgery”. The search strategy that was used for PubMed and other electronic databases is available on request from the corresponding author. In addition, the references of the selected articles were checked and the search was extended within cited references. Study selection and methodological screening was independently performed by two authors (GJR, JHB). In the case of disagreement, a third reviewer was consulted (AVN).

### Selection criteria

As case series are probably the most frequent type of surgical report in the literature (32), it was decided not to restrict the selection to a specific study design. As a consequence, studies were included if they used either within-group pre-post treatment comparisons or between-groups comparisons in a (randomized) controlled design.

In addition, studies were required to meet the following inclusion criteria: (i) investigating stroke and/or TBI in adults (irrespective of the phase of recovery); (ii) investigating the efficacy of surgical correction of equinovarus foot deformity (lengthening, release and/or transferring of muscles and/or tendons); (iii) being written as a full-length article in the English, German or Dutch languages and being published in a peer-reviewed journal between 1965 and March 2011. If two or more papers were published by the same group, and if (within these papers) aetiology of equinovarus foot deformity was comparable, only the study with the highest number of patients was included.

### Methodological quality assessment

The Oxford CEBM levels of evidence were used to grade the selected studies (33). The Index for Non-Randomized Studies (MINORS) was applied to further assess the quality of each study (34). Few validated instruments are available to assess the methodological quality of observational or non-randomized studies. MINORS is a validated list designed to assess the methodological quality of non-randomized (surgical) studies (either comparative or non-comparative) which comprises 12 items, of which the last 4 items apply only to comparative studies. Items are scored as 0 (not reported), 1 (reported but inadequate) or 2 (reported and adequate). The maximum score is 16 for non-comparative studies and 24 for comparative studies.

### Data extraction

Because no (randomized) controlled trials were identified, no pooling of data was allowed, either in a meta-analysis or in a best-evidence synthesis. The selected articles were analysed following the WHO ICF (available from: [www.who.int/icidh](http://www.who.int/icidh), 2001). The ICF is a classification of health and health-related domains at both individual and population level, commonly used within the field of rehabilitation medicine. It shifts the focus from cause to impact by describing how a disease can influence body structure and function, activity and participation.

## RESULTS

### Study selection

Fig. 1 shows the study selection process as a flow chart. The initial systematic search strategy in PubMed identified 320 relevant citations (on request available from the corresponding author). The search in the other databases did not yield additional articles. On the basis of the title, 228 studies were excluded. Another 46 studies were excluded based on their abstracts. Important reasons for exclusion were the use of interventions that did not fit within our definition, and the use of patient populations with an aetiology other than stroke or TBI. Full texts of the remaining 46 studies were examined. Screening the references of these studies revealed 4 additional articles (23, 35–37). From these 50 initially selected studies, 35 had to be excluded in second instance because they: (i) were




Fig. 1. Study selection process.

non-experimental (narrative reviews) ( $n=18$ ), (ii) included only children ( $n=3$ ), (iii) focused on equinus or equinovagis deformity ( $n=5$ ), (iv) evaluated outcome other than surgical efficacy ( $n=2$ ) (37, 38), (v) were published in a language other than English, Dutch or German ( $n=3$ ), or (vi) investigated cadavers ( $n=2$ ). Two other studies were excluded because they were part of larger studies published by the same group (39, 40). Ultimately, 15 studies met all inclusion criteria and were further analysed (35, 36, 41–53).

#### Methodological quality

Table I shows the methodological characteristics of the 15 studies included in the present review. Within our search limits, the earliest report regarding outcome of surgical correction of equinovarus foot deformity in patients with stroke or TBI dated from 1969 (45). All studies used a within-group design with pre-post treatment comparisons and were classified as case series (35, 36, 41, 43, 45, 47, 48, 50–53), case control (49), or historically controlled studies (42, 44, 46). Two studies (44, 49) used a prospective study design, whereas the other studies were conducted retrospectively. Only one study incorporated a group of healthy controls (49). All included studies were considered to be of the same, relatively low, level of methodological quality. They were scored as level 4 according to the definition of the Oxford Centre for Evidence-based Medicine. MINORS scores ranged from 4 to 14, emphasizing the methodological heterogeneity amongst the case-series designs (34).

#### Surgical techniques and comparisons

None of the studies compared surgical intervention with an alternative treatment. Three studies compared (uncontrolled, non-blinded) two techniques of surgical intervention; (i) Hosalkar et al. (42) differentiated two fixation techniques into the cuboid bone for split anterior tibialis tendon transfer (SPLATT), favouring the lateromedial over the dorsoplantar routing; (ii) Keenan et al. (44) compared toe flexor release with long toe flexor transfer to the calcaneus (to improve calf muscle strength), favouring the latter approach; and (iii) Morita et al. (46) compared long toe flexor transfer (to the fourth metatarsal bone on the dorsum of the foot) with whole anterior tibial tendon transfer (to the third cuneiform bone), favouring the first approach i.e. the use of long toe flexor transfer in dorsiflexion support. In one study, two groups were differentiated, based on (ab)normal electrical activity of the posterior tibial muscle in the stance phase, through which it was decided whether or not to lengthen this muscle. There were no differences in functional outcomes between both groups (49). In all but one study (53) the surgical procedures through which muscles and tendons were lengthened, released and/or transferred were described in detail.

#### Patient selection

The inclusion criteria in 13 studies were dominated by their retrospective nature. Patients were enrolled simply because they had had a surgical correction of equinovarus foot deformity. As a result, the original indication for this correction (i.e. the severity of the equinovarus foot deformity, additional impairments and/or the impact on activities of daily living) could not be properly traced in most studies. Some authors included only structural equinovarus deformity (36, 41), whereas others included only dynamic equinovarus deformity (47, 49). Four authors assigned surgery when extensive physiotherapy, nerve blocks and/or orthoses failed to correct the deviant equinovarus position (35, 45, 46, 49). In 4 studies regaining the ability to walk barefoot (i.e. without orthoses or orthopaedic footwear) was mentioned as a primary goal for surgery (46, 48, 51, 53). In 12 studies functional recovery of motor control after stroke (minimum of 6 months post-stroke) or TBI (minimum of 12 months post-TBI) was awaited before surgical correction was performed. In the remaining 3 studies information about the time since brain injury was lacking (35, 44, 45).

#### Gait assessment

The applied methods of gait assessment in the 15 studies are presented in Table II. Pre-operative instrumented gait analysis was used to assist in the surgical decision-making in 6 studies (41, 42, 44, 47, 49, 50), whereas 2 studies (36, 43) used only dynamic electromyography data for this purpose. The remaining 7 studies, all conducted before 1999, used qualitative (clinical) data in the pre-operative assessment. Only 3 studies used instrumented gait analysis in both the pre- and the postoperative assessments (41, 44, 49).

Table I. Methodological characteristics of the included studies

Authors	Study design	Minors score	Number of subjects	Age, years Mean [range]	Diagnosis	Time since injury, months Mean [range]	Intervention	Follow-up, months Mean [range]
Carda et al., 2009 (41)	Case series; uncontrolled; retrospective; single centre; non-blinded	13	177	50 [SD 14]	177 stroke	67 [no data]	135 a-l; 9 g-l; 32 gs-l; 174 fdl-r or fdl-l; 4 tp-l; 33 splatt; 71 ehl-t; 13 fh-l-t; 11 fdl-t	12 [constant]
Edwards et al., 1993 (36)	Case series; uncontrolled; retrospective; single centre; non-blinded	9	21 (24 feet)	41 [18-67]	9 stroke, 9 TBI, 2 other	34 [no data]	23 a-l + splatt + tf-r; 2 tp-l; 3 tp-r; 3 fp-r; 4 fh-l-t; (2 triple arthrodesis) I: 17 splatt; dorso plantar fixation	39 [12-79]
Hosalkar et al., 2008 (42)	Historically controlled; retrospective; single centre; non-blinded (comparing surgical techniques)	13	I: 17 II: 30	I: 46 [32-61] II: 42 [29-67]	47 TBI	18 [11-48]	II: 30 splatt; latero medial fixation note: additional procedures in both groups – not specified	I: 51 [37-84] II: 29 [24-48]
Keenan et al., 1984 (43)	Case series; uncontrolled; retrospective; single centre; non-blinded	10	54 (59 feet)	25 [14-50]	54 TBI	35 [SD 26]	53 a-l; 1 tp-l; 52 fdl-r; 34 fh-l-r; 59 splatt; 18 fh-l-t	49 [SD 23]
Keenan et al., 1999 (44)	Historically controlled; prospective; single centre; non-blinded (comparing surgical techniques)	14	I: 25 (31 feet) II: 30 (36 feet)	I: 40 [11-80] II: 36 [21-69]	I: 12 stroke, 11 TBI, 2 other II: 10 stroke, 17 TBI, 3 other	no data	I: 31 a-l + tf-r + splatt; 3 ehl-l; 13 tp-l	I: 41 [17-53]
Mooney et al., 1969 (45)	Case series; uncontrolled; retrospective; single centre; non-blinded	4	194	55 [17-84]	125 stroke	no data	II: 36 a-l + tf-r + fdl/fhl-t + splatt; 9 ehl-l; 15 tp-l note: additional proximal procedures in both groups	II: 24 [19-32]
Morita et al., 1998 (46)	Historically controlled; retrospective; single centre; non-blinded (comparing surgical techniques)	11	I: 110 II: 15	57 [32-78]	125 stroke	23 [6-132]	122 a-l; 122 tf-r; 82 splatt note: study includes description of proximal procedures I: 110 g-l + fdl-t + fh-l-t; 8 tp-l; 15 fdb-r + fhb-r II: 15 g-l + ta-t + fdl-r + fh-l-r; 3 fdb-r + fhb-r	33 [24-74]
Namdari et al., 2009 (47)	Case series; uncontrolled; retrospective; single centre; non-blinded	12	64	54 [24-74]	64 stroke	66 [17-523]	58 a-l; 30 tp-l; 7 ehl-l; 56 fp-r; 64 splatt; 61 fdl-t; 39 ehl-t	12 [3-42]
Ono et al., 1980 (48)	Case series; uncontrolled; retrospective; single centre; non-blinded	8	39	[18-76]	32 stroke, 1 TBI, 6 other	>12 [no data]	39 g-l + dl-t + fh-l-t note: in the same study 39 adult and 17 children	[46-98]



Table I. *Contd.*

Authors	Study design	Minors score	Number of subjects	Age, years Mean [range]	Diagnosis	Time since injury, months Mean [range]	Intervention	Follow-up, months Mean [range]
Pinzur et al., 1986 (49)	Case controlled; prospective; single centre; non-blinded (comparing surgical techniques)	13	I: 42 II: 12 (30 healthy controls)	stroke 57 [17–77] TBI 30 [18–51]	36 stroke, 16 TBI, 2 other	38 [12–204]	I: 42 a-l + fhl - r + fhb - r + fdl - r + ta II: 12 a-l + tp - l + fhl - l + fdl - r + ta	30 [24–62]
Reddy et al., 2008 (50)	Case series; uncontrolled; retrospective; single centre; non-blinded	12	26	55 [23–72]	26 stroke	75 [no data]	26 a-l + fhl - r + fdl - t + splatt; ? tp-l; ? ehl-t	18 [6–48]
Roper et al., 1978 (35)	Case series; uncontrolled; retrospective; single centre; non-blinded	6	50	43 [18–72]	stroke or TBI	no data	37 a-l; 13 tf-r; 8 tp-r; 23 splatt note: additional proximal procedures	no data
Tracy et al., 1976 (51)	Case series; uncontrolled; retrospective; single centre; non-blinded	6	35	40 [18–62]	22 stroke, 7 TBI, 6 other	36 [no data]	35 a-l + tp - l + tf - r + splatt	32 [4–73]
Vogt et al., 1998 (52)	Case series; uncontrolled; retrospective; single centre; non-blinded	12	69 (73 feet)	47 [8–79]	42 stroke, 15 TBI, 12 other	50 [no data]	49 a-l + fdl - r + fhl - r + splatt 9 a-l + splatt 6 fdl - r + fhl - r + splatt 9 splatt	44 [12–168]
Yamamoto et al., 1992 (53)	Case series; uncontrolled; retrospective; single centre; non-blinded	9	75	57 [no data]	75 stroke	18 [no data]	41 g-l + fdl - t + fhl - t 16 g-l + tf - r 3 g-l + tp - t 5 g-l + ta 5 g-l 5 tf - r	77 [no data]

SD: standard deviation; TBI: traumatic brain injury.

-l: lengthening; -r: release; -t: transfer; a: Achilles tendon; g: gastrocnemius and soleus; tf: toe flexor (intrinsic and extrinsic); fdl: flexor digitorum longus; fdb: flexor digitorum brevis; fhl: flexor hallucis longus; fhb: flexor hallucis brevis; tp: tibialis posterior ta: tibialis anterior (whole tendon); splatt: split anterior tibial tendon transfer; edl: extensor digitorum longus; ehl: extensor hallucis longus; fp: fascia plantaris.

### Outcomes

Outcome measures were categorized according to the ICF and are presented in Table II. In almost every study an improvement of the foot position following surgery was reported; however, only one study (41) statistically quantified this improvement. Walking speed was measured in 4 studies (41, 44, 46, 53), of which 2 reported a significant effect; in the study by Carda et al. (41) walking speed improved from 0.32 (m/s) prior to surgery to 0.40 m/s after surgery ( $p < 0.001$ ); Keenan et al. (44) found an improvement in walking speed from 0.36 m/s before to 0.5 m/s after surgery ( $p < 0.053$ ). Although improvement in walking speed appeared to be similar in two other studies (46, 53), the statistical significance of these findings was not supported. Both Carda et al. and Yamamoto et al. (41, 53) positively correlated improvement in walking speed with improvement in walking capacity. Again, only Carda et al. (41) statistically supported this correlation.

Orthotic use was reported in all 15 studies. In 14 studies the need for orthoses decreased post-operatively. In the study by Keenan et al. (43) the increase in post-operative orthotic use was explained by regaining walking capacity in the (formerly) non-ambulatory group. Three studies reported on the significance of decrease in orthotic use (44, 47, 52). Keenan et

al. (44) found a significant difference in the need of orthoses after surgery favouring the group in which the long toe flexors were transferred to the calcaneus ( $p = 0.025$ ).

Nine studies (36, 41–44, 46, 47, 50, 52) selected walking capacity as a primary outcome measure, whereas 4 studies did not report on walking capacity (48, 49, 51, 53) and 2 studies merely made a brief comment (35, 45). In these 9 studies, walking capacity was operationalized by ambulation categories ranging from non-ambulatory to unlimited community walker. Only 4 out of these 9 studies used a validated ambulation scale, i.e. the Walking Handicap Score (12) (41) and the Viosca score (54) (42, 47, 50). Although walking capacity seemed to improve in all studies, only 4 reported a statistically significant effect (41, 47, 50, 52). One other study suggested statistical significance, but no supporting data were presented (42).

In general, adverse effects and medical complications were rare, as were recurrences of foot deformities (Table III). Morita et al. (46) discussed the relatively high recurrence rates of hammer-toe and varus deformity in their study. When indicated, they now release the short toe flexors and lengthen the posterior tibial tendon, and report diminished recurrences of foot deformities.

Table II. Assessment and outcomes of surgical correction of equinovarus foot deformity in the included studies according to the International Classification of Functioning

Authors	Assessment of gait		Body structures and functions				Activities		Participation	
	Pre-	post-operative	Foot position		Walking speed	Spatiotemporal parameters	Orthotic use	Walking capacity	Patient satisfaction	Others
Carda et al., 2009 (41)	GA	GA	↑*	↑*	↑*	↑*	↓	↑*	NT	Propulsion ↑* Gait stability ↑*
Edwards et al., 1993 (36)	EMG	QA	↑	NT	NT	↑	Ø	↑	NT	
Hosalkar et al., 2008 (42)	I: GA	QA	↑	NT	NT	NT	↓	↑	NT	
	II: GA	QA	↑	NT	NT	NT	↓	↑	NT	
Keenan et al., 1984 (43)	EMG	QA	↑	NT	NT	NT	Ø	↑	NT	
Keenan et al., 1999 (44)	I: GA	QA	↑	↑	NT	NT	Ø	Ø	NT	
	II: GA	GA	↑	↑	↑*	↑	↓	↑	NT	
Mooney et al., 1969 (45)	QA	QA	↑	NT	NT	NT	Ø	↑	NT	
Morita et al., 1998 (46)	I: QA	QA/GA	↑	NT	↑	↑	↓	↑	NT	Propulsion ↑* Shift body weight ↑*
	II: QA	QA	↑	NT	NT	NT	↓	Ø	NT	
Namdari et al., 2009 (47)	GA	QA	↑	↑	NT	NT	↓*	↑*	NT	
Ono et al., 1980 (48)	QA	QA	↑	NT	NT	NT	Ø	NT	↑	
Pinzur et al., 1986 (49)	I: GA	GA	↑	↑	NT	↑	↓	NT	↑	Knee recurvation ↓
	II: GA	GA	↑	↑	NT	↑	↓	NT	↑	Knee recurvation ↓
Reddy et al., 2008 (50)	GA	QA	NT	NT	NT	NT	↓	↑*	NT	Use of non-operative therapy ↓ Knee recurvation ↓
Roper et al., 1978 (35)	QA	QA	↑	NT	NT	NT	↓	Ø	NT	
Tracy et al., 1976 (51)	QA	QA	↑	↑	NT	NT	↓	NT	NT	
Vogt, 1998 (52)	QA	QA	↑	↑	NT	NT	↓*	↑*	↑	
Yamamoto et al., 1992 (53)	QA	QA	NT	NT	↑	NT	↓	↑	↑	

↑\*: statistically significant improvement; ↓\* statistically significant deterioration; ↑: improvement in more than 50% of the patients (no statistical analysis); ↓: deterioration in more than 50% of the patients (no statistical analysis); Ø: no statistical difference or modification for less than 50% of the patients (no statistical analysis); NT: not tested; GA: instrumented gait analysis, including electromyography during walking; QA: qualitative assessment.

Table III. Adverse effects reported in the included studies

Author	Subjects <i>n</i>	Medical complications	Residual or recurrent deformities
Carda et al., 2009 (41)	177	3 deep vein thrombosis 4 wound infection 1 haematoma	3 toe curling
Edwards et al., 1993 (36)	21 (23 feet)	1 wound infection 1 tendon rupture	2 varus 1 toe curling
Hosalkar et al., 2008 (42)	I: 17 II: 30	3 screw pullout none	none none
Keenan et al., 1984 (43)	54 (59 feet)	1 wound infection with skin necrosis	none
Keenan et al., 1999 (44)	I: 25 (31 feet) II: 30 (36 feet)	no data no data	none 2 equinus
Mooney et al., 1969 (45)	194	6 wound infection 2 Achilles tendon rupture 1 death due to hepatitis	6 varus (15 valgus)
Morita et al., 1998 (46)	I: 110 II: 15	1 wound infection none	17 equinovarus, 21 toe curling 2 equinovarus, 6 toe curling
Namdari et al., 2009 (47)	64	no data	no data
Ono et al., 1980 (48)	39	no data	few toe curling
Pinzur et al., 1986 (49)	54	2 wound infection 2 tendon pullout	none
Reddy et al., 2008 (50)	26	none	no data
Roper et al., 1978 (35)	50	none	none
Tracy et al., 1976 (51)	35	1 chronic ulcer	3 equinovarus 3 toe curling
Vogt et al., 1998 (52)	69 (73 feet)	5 wound infection 5 tendon rupture 2 reflex dystrophy	3 equinovarus 6 toe curling
Yamamoto et al., 1992 (53)	75	no data	6 equinovarus 3 equinus 5 toe curling

## DISCUSSION

This systematic review was conducted to gain insight into the available evidence on the effects of soft-tissue surgery to correct equinovarus foot deformity following stroke or TBI. The rationale for this review was that, despite a paucity of well-conducted studies, clinical experience shows that surgical correction should be considered as a valuable treatment option. Several papers on the management of equinovarus foot deformity have been published (1–4, 55–57), but, as far as we are aware, there is no synthesis of experimental studies on surgical correction. The methodology of a systematic review offered the most objective approach (58).

The results of this review suggest that surgical correction of equinovarus foot deformity can be effective in re-obtaining a balanced foot position after stroke or TBI, which can improve walking capacity and diminish the need for an ankle-foot orthosis. However, the scientific evidence supporting surgi-

cal correction of equinovarus foot deformity is limited. All included studies were case series, scoring “level 4” evidence according to the definition of the Oxford Centre for Evidence-based Medicine (Grade C recommendation) (33). Indeed, the surgical procedures reported in this review have all been evaluated in non-controlled studies. In general, it is estimated that randomized controlled trials (RCTs) account for less than 10% of the evidence base for surgical interventions (59). Nevertheless, despite the inherent difficulties in conducting RCTs, well-controlled studies are also essential in proving the effects of surgical procedures, for instance by using a control group that will receive the studied intervention in a later stage. In the present review, all but two of the included studies were retrospective, which implies that the data had been collected before stating an hypothesis in the context of a scientific study. As a consequence these data are highly subjective to information bias, which compromises the internal validity of the studies. On the other hand, in observational studies such as case series, the investigator does not purposely control which patients are selected and what intervention they receive, which implies that the results are probably closer to routine clinical practice and, therefore, have a relatively high external validity. In addition, depending on their methodological quality, observational studies can make valid statements about the safety of surgical intervention (60). All studies in this review showed that adverse effects and medical complications were rare, which makes it fair to conclude that surgical correction of equinovarus foot deformity following stroke or TBI can be safely performed.

An adequate *selection* of patients suitable for surgical equinovarus foot correction after stroke or TBI should be the first challenge in future research. Due to methodological shortcomings and poor description, criteria for selecting the most suitable patients cannot be derived from the present review. In current rehabilitation practice, surgical correction often seems to be an afterthought in treatment options. Patients with a structural equinovarus deformity or those with severe spasticity who failed to respond to alternative treatment (such as an orthosis or chemodenervation) will eventually be considered eligible for surgery. Surgical correction might, however, be considered in an earlier stage. In addition, based on this review, patients with dynamic equinovarus foot deformity should also be considered for surgical intervention. Indeed, one of the major advantages of surgical correction of equinovarus foot deformity is that patients regain the opportunity to walk barefoot, which is essential when getting out of bed during the night, when taking a shower, or during leisure activities. During such activities, many patients wish to be more independent of, for example, orthosis, which is often neglected by rehabilitation professionals. The advantage of walking barefoot favours surgical correction over orthotics, even if the effectiveness of these different interventions on walking capacity and gait velocity would be comparable. This review showed that recurrences of foot deformity after surgery are rare, which is equally important because it indicates that in many patients surgical correction is not only a safe, but also a permanent,



treatment option. Another aspect of patient *selection* is the timing of surgery post-stroke or post-TBI. Neurological recovery is believed to continue up to 6–9 months after stroke and up to 12–18 months after TBI, which is why surgical correction of equinovarus foot deformity is often postponed until after this recovery period (see Table I). However, the benefits or consequences of postponing surgery have never been formally investigated. Recent evidence, showing that within 3 weeks post-stroke muscle activation patterns in the affected leg are more or less established, seems to support the notion of earlier surgical intervention post-stroke, which may prevent unnecessary development of compensatory gait strategies (61). Of course, it should be kept in mind that surgical correction of equinovarus foot deformity is an elective procedure. Patients with poor vascularization, or other conditions that pose risks to the operation should be excluded.

The second challenge for future research is standardization of *preoperative planning*. From the selected studies a tendency towards thorough (instrumented) gait assessment prior to surgical planning can be determined (Table II). Developing evidence-based guidelines regarding the surgical correction of equinovarus foot deformity will depend on a structured approach. Equinovarus foot deformity can result from a variety of imbalanced muscle forces for which a single, best operation is not available. Thus, surgical correction should, pre-eminently, be patient tailored. Instrumented gait analysis offers the possibility to collect objective and repeatable data on the severity and dynamics of the equinovarus deformity and its consequences for walking capacity (62). Gait analysis, at present, is probably the most powerful instrument to assist clinicians in surgical decision-making and to assess the outcome of surgery (63, 64). Because the activation pattern of most operated muscles alters very little (38), pre-operative gait analysis (including electromyography during gait) is essential in determining the appropriate surgical plan, since it is a good indication of the type of muscle activity to expect post-operatively. Fuller et al. (37) demonstrated that instrumented gait analysis altered the surgical planning for patients with equinovarus foot deformity and produced a higher agreement among surgeons. Introducing instrumented gait analysis as a gold standard not only enables clinicians (and researchers) to exchange reliable and detailed information about patients, but also creates the opportunity to (remotely) consult colleagues or other experts to determine the suitability of an individual patient for a specific surgical procedure (65).

The third challenge concerns uniformity in *outcome measures*. Preferably, all domains of the ICF (functions and structures, activities and participation) should be assessed (Table II). Outcome measures should evaluate the desired treatment effect, but also allow for comparing effectiveness of treatment alternatives. The psychometric properties (reliability, validity, responsiveness) of outcome measures are important criteria for selection (66–68). Instrumented gait analysis produces reliable and repeatable data on spatiotemporal, kinetic and kinematic gait characteristics. By objectively quantifying underlying impairments, gait analysis is not only essential in preoperative planning, but also in evaluating the effect of surgical correc-

tion on the deviant ankle-foot position and on the intended improvement of walking capacity. Furthermore, gait analysis allows for detailed comparison with treatment alternatives, such as ankle-foot orthotics, chemodenervation and functional electrical stimulation (63, 69, 70). Of the included studies, only 3 used instrumented gait analysis in both the pre- and post-operative assessment (Table II).

As equinovarus deformity compromises (barefooted) walking, regaining walking capacity will be the first objective of all interventions that aim to correct equinovarus deformity, including surgical correction. Therefore, outcome measures should at least incorporate validated measures of walking capacity. Such measures were used by only 4 studies included in this review. Based on their psychometric properties, the Six-Minute Walk Test (6MWT) (71) and the Functional Ambulation Categories (FAC) (72, 73) appear to be suitable to measure walking capacity. Walking speed can additionally be regarded as a simple and valid parameter of gait, since it is highly correlated with walking capacity (74). Walking speed can, for instance, be measured accurately with the Ten-Meter Comfortable Walk Test (10MCWT) (75). It should be acknowledged, however, that these capacity measures only provide information about what patients *are able to do*, but not about their actual performance (i.e. activities) in daily life. In addition, most clinical outcome measures are unable to quantify the quality of performance while executing meaningful tasks, such as gait. In this perspective, it seems essential to assess the total ambulation time over a representative time period. In addition, validated questionnaires on societal participation will give insight into which activities patients are involved in and provide a more or less complete picture of the benefits of the (surgical) intervention as experienced by individual patients.

A second major objective of surgical intervention is to eliminate (or at least reduce) the need for an ankle-foot orthosis for walking. All studies included in this review reported on the pre- and post-treatment need for orthoses. None of these studies, however, quantitatively compared the effect of an ankle-foot orthosis with that of the surgical correction. In our opinion, recommending a surgical correction would be appropriate only if the expected effect on walking capacity is at least comparable to the effect of an orthosis.

In conclusion, despite the methodological shortcomings of the included studies, it seems fair to state that surgical correction of equinovarus foot deformity following stroke or TBI is a safe and more or less “permanent” treatment option with a good chance of improving walking capacity and of diminishing the need for orthotic use. However, the level of evidence of the selected studies was low and used outcome measures were heterogeneous. It was not possible to conduct a quantitative analysis (i.e. meta-analysis) of the collected results. Further validation of surgical correction of equinovarus foot deformity following stroke or TBI is required, using higher-level study designs (prospective cohort, RCT) with validated assessment tools. Studies that incorporate control groups are necessary to compare surgical techniques with other interventions, with the aim of generating evidence upon which treatment algorithms can be based.

## REFERENCES

1. Deltombe T, Gustin T, De Cloedt P, Vandemeulebroecke M, Hanson P. The treatment of spastic equinovarus foot after stroke. *Crit Rev Phys Rehabil Med* 2007; 19: 195–211.
2. Forese L, Wootten M, Kadaba MP, McCann PD. Surgical management of equinovarus deformity in the adult with head injury. *Orthop Rev* 1993; 22: 1001–1010.
3. Roper BA. The orthopedic management of the stroke patient. *Clin Orthop Relat Res* 1987; 219: 78–86.
4. Botte MJ, Bruffey JD, Copp SN, Colwell CW. Surgical reconstruction of acquired spastic foot and ankle deformity. *Foot Ankle Clin* 2000; 5: 381–416.
5. Dobkin BH. Clinical practice. Rehabilitation after stroke. *N Engl J Med* 2005; 352: 1677–1684.
6. Katz DI, White DK, Alexander MP, Klein RB. Recovery of ambulation after traumatic brain injury. *Arch Phys Med Rehabil* 2004; 85: 865–869.
7. Jorgensen HS, Nakayama H, Raaschou HO, Olsen TS. Recovery of walking function in stroke patients: the Copenhagen Stroke Study. *Arch Phys Med Rehabil* 1995; 76: 27–32.
8. Kollen B, van de Port I, Lindeman E, Twisk J, Kwakkel G. Predicting improvement in gait after stroke: a longitudinal prospective study. *Stroke* 2005; 36: 2676–2680.
9. Veerbeek JM, Van Wegen EE, Harmeling-Van der Wel BC, Kwakkel G, EPOS Investigators. Is accurate prediction of gait in nonambulatory stroke patients possible within 72 hours poststroke? The EPOS study. *Neurorehabil Neural Repair* 2011; 25: 268–274.
10. Heiden JS, Small R, Caton W. Severe head injury and outcome; a prospective study. In: Popp R, Bourke RS, Nelson LR, editors. *Neural trauma*. New York: Raven Press; 1979.
11. Brandstater ME, de Bruin H, Gowland C, Clark BM. Hemiplegic gait: analysis of temporal variables. *Arch Phys Med Rehabil* 1983; 64: 583–587.
12. Perry J, Garrett M, Gronley JK, Mulroy SJ. Classification of walking handicap in the stroke population. *Stroke* 1995; 26: 982–989.
13. Lord SE, McPherson K, McNaughton HK, Rochester L, Weatherall M. Community ambulation after stroke: how important and obtainable is it and what measures appear predictive? *Arch Phys Med Rehabil* 2004; 85: 234–239.
14. Gage JR. Gait analysis. An essential tool in the treatment of cerebral palsy. *Clin Orthop Relat Res* 1993; 288: 126–134.
15. Weerdesteyn V, de Niet M, van Duijnhoven HJ, Geurts AC. Falls in individuals with stroke. *J Rehabil Res Dev* 2008; 45: 1195–1213.
16. Malas B, Kacen M. Orthotic management in patients with stroke. *Top Stroke Rehabil* 2001; 7: 38–45.
17. Pittcock SJ, Moore AP, Hardiman O, Ehler E, Kovac M, Bojakowski J, et al. A double-blind randomised placebo-controlled evaluation of three doses of botulinum toxin type A (Dysport) in the treatment of spastic equinovarus deformity after stroke. *Cerebrovasc Dis* 2003; 15: 289–300.
18. Kocabas H, Salli A, Demir AH, Ozerbil OM. Comparison of phenol and alcohol neurolysis of tibial nerve motor branches to the gastrocnemius muscle for treatment of spastic foot after stroke: a randomized controlled pilot study. *Eur J Phys Rehabil Med* 2010; 46: 5–10.
19. Foley N, Murie-Fernandez M, Speechley M, Salter K, Sequeira K, Teasell R. Does the treatment of spastic equinovarus deformity following stroke with botulinum toxin increase gait velocity? A systematic review and meta-analysis. *Eur J Neurol* 2010; 17: 1419–1427.
20. Bollens B, Deltombe T, Detrembleur C, Gustin T, Stoquart G, Lejeune TM. Effects of selective tibial nerve neurotomy as a treatment for adults presenting with spastic equinovarus foot: a systematic review. *J Rehabil Med* 2011; 43: 277–282.
21. Kottink AIR, Oostendorp LJM, Buurke JH, Nene AV, Hermens HJ, IJzerman MJ. The orthotic effect of functional electrical stimulation on the improvement of walking in stroke patients with a dropped foot: a systematic review. *Artif Organs* 2006; 28: 577–586.
22. IJzerman MJ, Renzenbrink GJ, Geurts AC. Neuromuscular stimulation after stroke: from technology to clinical deployment. *Expert Rev Neurother* 2009; 9: 541–552.
23. Kamath AF, Pandya MD, Namdari S, Hosalkar H, Keenan MA. Surgical technique for the correction of adult spastic equinovarus foot. *Tech Foot Ankle* 2009; 8: 160–167.
24. Keenan MA. The management of spastic equinovarus deformity following stroke and head injury. *Foot Ankle Clin* 2011; 16: 499–514.
25. Australian Clinical Guidelines for Stroke Management. Clinical Guidelines for Stroke Rehabilitation and Recovery [Internet]. 2005. Available from: [www.strokefoundation.com.au](http://www.strokefoundation.com.au).
26. Teasell R, Foley N, Salter K. 13th edition of Evidence Based Review of Stroke Rehabilitation (funding of the Canadian stroke network) [Internet]. 2010. Available from: [www.ebrsr.com](http://www.ebrsr.com).
27. International Society of Prosthetics and Orthotics. Report of a consensus conference on orthotic management of stroke patients. 2003. Available from: <http://www.ispoint.org>.
28. van Til JA, Renzenbrink GJ, Dolan JG, IJzerman MJ. The use of the analytic hierarchy process to aid decision making in acquired equinovarus deformity. *Arch Phys Med Rehabil* 2008; 89: 457–462.
29. Stroke – 1989. Recommendations on stroke prevention, diagnosis, and therapy. Report of the WHO Task Force on Stroke and other Cerebrovascular Disorders. *Stroke* 1989; 20: 1407–1431.
30. The Traumatic Brain Injury Model System of Care. Funded by the National Institute on Disability and Rehabilitation Research. Available from: <http://www.tbindsc.org>.
31. WHO. International Classification of Functioning, Disability and Health: ICF. Geneva: World Health Organization; 2001.
32. Ridgway PF, Guller U. Interpreting study designs in surgical research: a practical guide for surgeons and surgical residents. *J Am Coll Surg* 2009; 208: 635–645.
33. OCEBM Levels of Evidence Working Group. “The Oxford 2011 Levels of Evidence”. Oxford Centre for Evidence-Based Medicine [Internet]. Available from: <http://www.cebm.net/index.aspx?o=5653>.
34. Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ J Surg* 2003; 73: 712–716.
35. Roper BA, Williams A, King JB. The surgical treatment of equinovarus deformity in adults with spasticity. *J Bone Joint Surg Br* 1978; 60-B: 533–535.
36. Edwards P, Hsu J. SPLATT combined with tendo achilles lengthening for spastic equinovarus in adults: results and predictors of surgical outcome. *Foot Ankle* 1993; 14: 335–338.
37. Fuller DA, Keenan MA, Esquenazi A, Whyte J, Mayer NH, Fidler-Sheppard R. The impact of instrumented gait analysis on surgical planning: treatment of spastic equinovarus deformity of the foot and ankle. *Foot Ankle Int* 2002; 23: 738–743.
38. Waters RL, Frazier J, Garland DE, Jordan C, Perry J. Electromyographic gait analysis before and after operative treatment for hemiplegic equinus and equinovarus deformity. *J Bone Joint Surg Am* 1982; 64: 284–288.
39. Morita S, Yamamoto H, Furuya K. Anterior transfer of the toe flexors for equinovarus deformity due to hemiplegia. *J Bone Joint Surg Br* 1994; 76: 447–449.
40. Carda S, Molteni F, Bertoni M, Zerbinati P, Invernizzi M, Cisari C. Extensor hallucis longus transfer as an alternative to split transfer of the tibialis anterior tendon to correct equinovarus foot in hemiplegic patients without overactivity of tibialis anterior. *J Bone Joint Surg Br* 2010; 92: 1262–1266.
41. Carda S, Bertoni M, Zerbinati P, Rossini M, Magoni L, Molteni F. Gait changes after tendon functional surgery for equinovarus foot in patients with stroke: assessment of temporo-spatial, kinetic, and kinematic parameters in 177 patients. *Am J Phys Med Rehabil* 2009; 88: 292–301.

42. Hosalkar H, Goebel J, Reddy S, Pandya NK, Keenan MA. Fixation techniques for split anterior tibialis transfer in spastic equinovarus feet. *Clin Orthop Relat Res* 2008; 466: 2500–2506.
43. Keenan MA, Creighton J, Garland DE, Moore T. Surgical correction of spastic equinovarus deformity in the adult head trauma patient. *Foot Ankle* 1984; 5: 35–41.
44. Keenan MA, Lee GA, Tuckman AS, Esquenazi A. Improving calf muscle strength in patients with spastic equinovarus deformity by transfer of the long toe flexors to the Os calcis. *J Head Trauma Rehabil* 1999; 14: 163–175.
45. Mooney V, Goodman F. Surgical approaches to lower-extremity disability secondary to strokes. *Clin Orthop Relat Res* 1969; 63: 142–152.
46. Morita S, Muneta T, Yamamoto H, Shinomiya K. Tendon transfer for equinovarus deformed foot caused by cerebrovascular disease. *Clin Orthop Relat Res* 1998; 350: 166–173.
47. Namdari S, Park MJ, Baldwin K, Hosalkar HS, Keenan MA. Effect of age, sex, and timing on correction of spastic equinovarus following cerebrovascular accident. *Foot Ankle Int* 2009; 30: 923–927.
48. Ono K, Hiroshima K, Tada K, Inoue A. Anterior transfer of the toe flexors for equinovarus deformity of the foot. *Int Orthop* 1980; 4: 225–229.
49. Pinzur MS, Sherman R, DiMonte-Levine P, Kett N, Trimble J. Adult-onset hemiplegia: changes in gait after muscle-balancing procedures to correct the equinus deformity. *J Bone Joint Surg Am* 1986; 68: 1249–1257.
50. Reddy S, Kusuma S, Hosalkar H, Keenan MA. Surgery can reduce the nonoperative care associated with an equinovarus foot deformity. *Clin Orthop Relat Res* 2008; 466: 1683–1687.
51. Tracy HW. Operative treatment of the plantar-flexed inverted foot in adult hemiplegia. *J Bone Joint Surg Am* 1976; 58: 1142–1145.
52. Vogt JC. Split anterior tibial transfer for spastic equinovarus foot deformity: retrospective study of 73 operated feet. *J Foot Ankle Surg* 1998; 37: 2–7.
53. Yamamoto H, Okumura S, Morita S, Obata K, Furuya K. Surgical correction of foot deformities after stroke. *Clin Orthop Relat Res* 1992; 282: 213–218.
54. Viosca E, Martinez JL, Almagro PL, Gracia A, Gonzalez C. Proposal and validation of a new functional ambulation classification scale for clinical use. *Arch Phys Med Rehabil* 2005; 86: 1234–1238.
55. Asencio G, Pelissier J, Privat JM, Bertin R, Megy B, Leonardi C. Surgery of the foot in equinus deformity in hemiplegic adults. *Rev Chir Orthop Reparatrice Appar Mot* 1993; 79: 41–48.
56. Jaivin JS, Bishop JO, Braly WG, Tullos HS. Management of acquired adult dropfoot. *Foot Ankle* 1992; 13: 98–104.
57. Lawrence SJ, Botte MJ. Management of the adult, spastic, equinovarus foot deformity. *Foot Ankle Int* 1994; 15: 340–346.
58. Cook DJ, Mulrow CD, Haynes RB. Systematic reviews: synthesis of best evidence for clinical decisions. *Ann Intern Med* 1997; 126: 376–380.
59. McCulloch P, Taylor I, Sasako M, Lovett B, Griffin D. Randomised trials in surgery: problems and possible solutions. *BMJ* 2002; 324: 1448–1451.
60. Kooistra B, Dijkman B, Einhorn TA, Bhandari M. How to design a good case series. *J Bone Joint Surg Am* 2009; 91 Suppl 3: 21–26.
61. Buurke JH, Nene AV, Kwakkel G, Erren-Wolters V, Ijzerman MJ, Hermens HJ. Recovery of gait after stroke: what changes? *Neurorehabil Neural Repair* 2008; 22: 676–683.
62. Yavuzer G, Oken O, Elhan A, Stam HJ. Repeatability of lower limb three-dimensional kinematics in patients with stroke. *Gait Posture* 2008; 27: 31–35.
63. Patrick JH, Keenan MA. Gait analysis to assist walking after stroke. *Lancet* 2007; 369: 256–257.
64. Wren TA, Otsuka NY, Bowen RE, Scaduto AA, Chan LS, Sheng M, et al. Influence of gait analysis on decision-making for lower extremity orthopaedic surgery: Baseline data from a randomized controlled trial. *Gait Posture* 2011; 34: 364–369.
65. Buurke JH, Kleissen RF, Nene A, Bloo JK, Renzenbrink GJ, Zeegers AV, et al. A feasibility study of remote consultation to determine suitability for surgery in stroke rehabilitation. *J Telemed Telecare* 2004; 10: 108–112.
66. Salter K, Jutai JW, Teasell R, Foley NC, Bitensky J. Issues for selection of outcome measures in stroke rehabilitation: ICF Body Functions. *Disabil Rehabil* 2005; 27: 191–207.
67. Salter K, Jutai JW, Teasell R, Foley NC, Bitensky J, Bayley M. Issues for selection of outcome measures in stroke rehabilitation: ICF Activity. *Disabil Rehabil* 2005; 27: 315–340.
68. Salter K, Jutai JW, Teasell R, Foley NC, Bitensky J, Bayley M. Issues for selection of outcome measures in stroke rehabilitation: ICF Participation. *Disabil Rehabil* 2005; 27: 507–528.
69. van Swigchem R, Vloothuis J, den Boer J, Weerdesteyn V, Geurts AC. Is transcutaneous peroneal stimulation beneficial to patients with chronic stroke using an ankle-foot orthosis? A within-subjects study of patients' satisfaction, walking speed and physical activity level. *J Rehabil Med* 2010; 42: 117–121.
70. Caty GD, Detrembleur C, Bleyenheuft C, Lejeune TM. Reliability of lower limb kinematics, mechanics and energetics during gait in patients after stroke. *J Rehabil Med* 2009; 41: 588–590.
71. Fulk GD, Echternach JL, Nof L, O'Sullivan S. Clinometric properties of the six-minute walk test in individuals undergoing rehabilitation poststroke. *Physiother Theory Pract* 2008; 24: 195–204.
72. Mehrholz J, Wagner K, Rutte K, Meissner D, Pohl M. Predictive validity and responsiveness of the functional ambulation category in hemiparetic patients after stroke. *Arch Phys Med Rehabil* 2007; 88: 1314–1319.
73. Holden MK, Gill KM, Magliozzi MR. Gait assessment for neurologically impaired patients. Standards for outcome assessment. *Phys Ther* 1986; 66: 1530–1539.
74. Schmid A, Duncan PW, Studenski S, Lai SM, Richards L, Perera S, et al. Improvements in speed-based gait classifications are meaningful. *Stroke* 2007; 38: 2096–2100.
75. Flansbjerg UB, Holmback AM, Downham D, Patten C, Lexell J. Reliability of gait performance tests in men and women with hemiparesis after stroke. *J Rehabil Med* 2005; 37: 75–82.